

1 Citation: **Hirt C** (2011) Assessment of EGM2008 over Germany using accurate quasigeoid heights from vertical deflections,  
2 GCG05 and GPS/levelling. *Zeitschrift für Geodäsie, Geoinformation und Landmanagement (zfv)* 136(3): 138-149.

# 3 **Assessment of EGM2008 over Germany using accurate** 4 **quasigeoid heights from vertical deflections, GCG05 and** 5 **GPS/levelling**

6

7 **Christian Hirt**

8 Western Australian Centre for Geodesy & The Institute for Geoscience Research

9 Curtin University of Technology, GPO Box U1987, Perth, WA 6845, Australia

10 Fax: +61 8 9266 2703; Email: [c.hirt@curtin.edu.au](mailto:c.hirt@curtin.edu.au)

11

## 12 **Summary**

13 EGM2008 is a high-resolution global model of Earth's gravity field that allows computation  
14 of quasigeoid heights and further functionals down to a resolution of 5 arc minutes. The  
15 present paper assesses EGM2008 over Germany by means of quasigeoid heights from the  
16 German GCG05 quasigeoid model and GPS/levelling points, and quasigeoid height  
17 differences from astronomical levelling. Residual terrain model (RTM) data is used for the  
18 computation of RTM quasigeoid heights, serving to augment the resolution of EGM2008 at  
19 scales shorter than 5 arc minutes. For quasigeoid heights, the comparisons show a RMS (root  
20 mean square) agreement of ~3 cm between EGM2008 and GCG05 as well as EGM2008 and  
21 GPS/levelling. The residuals between EGM2008 (augmented with RTM) and astrogeodetic  
22 quasigeoid height differences are near or at the cm-level for two local test areas. The  
23 comparisons show the very good quality of EGM2008 over Germany, which serves as an  
24 example region where dense gravity sets were used for the model's development.

## 25 **Zusammenfassung**

26 EGM2008 ist ein hochauflösendes globales Erdschwerefeldmodell, das zur Berechnung von  
27 Quasigeoidhöhen und anderen Schwerefeldfunktionalen mit einer Auflösung von 5  
28 Bogenminuten verwendet werden kann. Der vorliegende Beitrag bewertet EGM2008 mit  
29 Hilfe des Quasigeoidmodells GCG05, einem GPS/Nivellement Datensatz und  
30 astrogeodätisch bestimmten Differenzen von Quasigeoidhöhen. Residuale Geländedaten  
31 (RTM) werden zur Berechnung von RTM Quasigeoidhöhen genutzt, die EGM2008 auf  
32 Skalen kürzer als 5 Bogenminuten ergänzen. Die Vergleiche zeigen mittlere quadratische  
33 Abweichungen (RMS) von etwa 3 cm zwischen EGM2008 und GCG05 bzw. den  
34 GPS/Nivellementspunkten. Die Residuen zwischen EGM2008 und astrogeodätischen  
35 Differenzen von Quasigeoidhöhen in zwei lokalen Testgebieten sind auf oder nahe dem cm-  
36 Niveau. Der Beitrag zeigt die sehr gute Qualität von EGM2008 über Deutschland als Beispiel  
37 für Gebiete, in denen umfangreiche Schwerewerte für die EGM2008 Modellierung verwendet  
38 wurden.

39 **Keywords:** EGM2008, quasigeoid, GCG05, GPS/levelling, astronomical levelling, residual  
40 terrain modelling (RTM)

## 41 **1. Introduction**

42 With the computation and release of the Earth Gravitational Model EGM2008 (Pavlis et al.  
43 2008) in April 2008, a major advancement was made in high-resolution global gravity field  
44 modelling. Developed by the U.S. National Geospatial Agency (NGA), EGM2008 is the first-  
45 ever global model that is capable of resolving the Earth's gravity field beyond spherical  
46 harmonic degree 2000. The EGM2008 set of spherical harmonic coefficients is complete to  
47 degree 2190 and order 2159. It allows computation of various gravity field functionals – such  
48 as quasigeoid heights, gravity anomalies and vertical deflections – globally with a spatial  
49 resolution of ~5 arc minutes, or ~9 km in the latitudinal direction. EGM2008 is freely  
50 available from <http://earth-info.nga.mil/GandG/wgs84/gravitymod/egm2008/index.html>.

51 A number of external evaluation studies on EGM2008 have already been carried out using  
52 'ground-truth' gravity field observations over several countries (Newton's Bulletin 2009).  
53 The comparisons made in the 25 studies presented in Newton's Bulletin (2009) provide  
54 evidence of low EGM2008 commission errors, i.e., the uncertainties of EGM2008-derived  
55 functionals, particularly over areas where EGM2008 is based on dense gravity data sets.

56 As a consequence of its high spatial resolution and accuracy, EGM2008 represents a large  
57 part of the gravity field spectrum. Because the residual gravity field signals are very small,  
58 EGM2008-based regional gravity field modelling encounters new challenges (e.g.,  
59 Featherstone et al. 2010). Recent examples of regional gravity field modelling using  
60 EGM2008 are given by Featherstone et al. (2010) for Australia, Claessens et al. (2011) for  
61 New Zealand, Roman et al. (2010) for the United States and Denker et al. (2009) and Ihde et  
62 al. (2010) for Europe.

63 Beyond its resolution, that is at scales finer than ~5 arc minutes, EGM2008 is not capable of  
64 representing the high-frequency constituents of Earth's gravity field. The neglect of high-  
65 frequency content by a harmonic model like EGM2008 is known as omission error (Torge  
66 2001 p. 273; Gruber 2009). For quasigeoid heights derived from EGM2008, Jekeli et al.  
67 (2009) estimated the EGM2008 omission error to be ~ 4 cm. This is a global estimate which  
68 may vary for different types of terrain. Little or no attempt was made to model and account  
69 for the EGM2008-omitted high-frequency signals in the evaluation reports on EGM2008  
70 (Newton's Bulletin 2009).

71 To model and reduce the omission error, one common strategy is the remove-compute-restore  
72 (RCR) approach, where the fine-structure is sourced from residual gravity (Torge 2001, p.  
73 286). As is known, many regional geoid or quasigeoid models are based on this method,  
74 including the above mentioned regional models based on EGM2008. Alternatively, residual  
75 terrain model (RTM) data (Forsberg 1984) can be used in elevated terrain as a source to  
76 recover parts of the omission error (Hirt 2010, Hirt et al. 2010a, 2010b). Not only is RTM-  
77 based omission error modelling advantageous for accurate prediction of functionals (e.g., Hirt

78 2010), but it also facilitates the assessment of EGM2008 (and other spherical harmonic  
79 models) with ground-truth observations (Hirt et al. 2010a).

80 The present paper assesses EGM2008 (Section 2) over Germany using the RTM  
81 augmentation technique (Section 3) and three different sources of accurate quasigeoid heights  
82 (Section 4). These are (i) the German Combined Quasigeoid GCG05 (Liebsch et al. 2006,  
83 Schirmer et al. 2006), (ii) a set of quasigeoid heights from GPS/levelling points (Ihde and  
84 Sacher 2002) and (iii) two local profiles of astrogeodetic quasigeoid differences (Hirt et al.  
85 2008, Hirt and Flury 2008). Section 5 then presents and discusses the results of the  
86 comparisons with EGM2008. A first focus is placed on the inclusion of omission error  
87 estimates from RTM data, so as to ‘bridge’, to some extent, the spectral gap between the  
88 EGM2008 quasigeoid heights and the comparison data (Hirt et al. 2010b). A second focus is  
89 set on the role of the station heights at which EGM2008 is evaluated (Section 5). Germany  
90 was selected not only because of the sufficiently accurate comparison data sets available, but  
91 also as an example region where dense gravity data sets were available and used for the  
92 EGM2008 model construction (see Pavlis et al. 2008).

93 This paper is complementary to other studies comparing EGM2008 against terrestrial data  
94 sets over Germany. For example, Förste et al. (2009) and Gruber (2009) used GPS/levelling  
95 points to evaluate EGM2008 in comparison to other geopotential models with focus on the  
96 long- and medium-wavelength domain. A study by Ihde et al. (2010) used GPS/levelling  
97 points for EGM2008 evaluation while results from a comparison between astrogeodetic  
98 quasigeoid height differences and EGM2008 were reported in Berichte (2010). However, the  
99 EGM2008 omission error beyond its maximum degree of expansion was neither modelled  
100 nor reduced in these studies. For comparisons among vertical deflections and EGM2008 over  
101 Germany, see Voigt et al. (2008), Ihde et al. (2010), Hirt (2010) and Hirt et al. (2010a).

## 102 **2. EGM2008**

103 A paper outlining the details of EGM2008 has not yet become available, however a general  
104 overview of EGM2008 is given in Pavlis et al. (2008) with background information on the  
105 model’s development presented in Kenyon et al. (2007), Pavlis et al. (2007), Holmes et al.  
106 (2007), Pavlis and Saleh (2004), Pavlis et al. (2004) and Saleh and Pavlis (2003). EGM2008  
107 consists of a total of ~4.8 million spherical harmonic coefficients complete to degree and  
108 order 2159, with additional spherical harmonic coefficients to degree 2190 and order 2159  
109 (EGM Development Team 2008). The EGM2008 geopotential model is available free-of-  
110 charge, together with accompanying products such as a spherical harmonic model of Earth’s  
111 topography, grids of commission error estimates for different gravity field functionals and a  
112 high-degree synthesis software.

113 EGM2008 is based on the GRACE (Gravity Recovery and Climate Experiment)-only gravity  
114 field model ITG-GRACE03S (Mayer-Gürr 2007) which provides a highly-accurate  
115 description of the long- and medium-wavelength gravity field spectrum up to degree and  
116 order 180. The ITG-GRACE03S model incorporates almost 6 years of GRACE gravity field  
117 observations. The second input data set is a global grid of 5’×5’ area-mean gravity anomalies


118 (band-limited to degree 2160) that was constructed from high-resolution topographic data  
119 (Pavlis et al. 2007, Pavlis and Saleh 2004), altimetry-derived gravity over the oceans (e.g.,  
120 Andersen et al. 2010), and other sources of gravity data, particularly point gravity  
121 measurements (Pavlis et al. 2008).

122 The global grid of surface area-mean gravity anomalies was harmonically analysed to derive  
123 a set of ellipsoidal harmonic coefficients (Pavlis et al. 2004, Holmes and Pavlis 2007). The  
124 ITG-GRACE03S satellite gravity model was converted from spherical to ellipsoidal  
125 harmonics by means of Jekeli's (1988) transformation. The ellipsoidal harmonic coefficients  
126 of both input data sets were then combined through a least-squares adjustment procedure  
127 (Pavlis et al. 2008). The resulting ellipsoidal harmonic spectrum (complete to degree and  
128 order 2159) was finally back-converted to spherical harmonics using Jekeli's (1988)  
129 algorithm. Because this transformation preserves the maximum order, but not the maximum  
130 degree of the harmonic series expansion (see also Holmes and Pavlis 2007), some additional  
131 coefficients (to degree 2190 and order 2159) occur in spherical harmonic representation. It is  
132 recommended not to neglect these additional coefficients (cf. Holmes and Pavlis 2007).  
133 Hence, the EGM2008 spherical harmonic coefficients should be expanded to degree 2190  
134 rather than only to 2159 or 2160 when being employed in practical applications.

135 The very good quality of EGM2008 in the long and medium wavelengths is mainly due to  
136 using GRACE satellite gravity field observations, supported by the spectral content implied  
137 in this band by terrestrial gravity anomalies. EGM2008's spectral band between 181 to 2159  
138 (in terms of ellipsoidal harmonics) originates solely from the 5'×5' area-mean gravity  
139 anomalies (see above). Because of the inhomogeneous and incomplete global coverage by  
140 surface gravity observations, the NGA 5'×5' area-mean gravity data base is of varying quality  
141 (Pavlis et al. 2008, pp. 2-4). As a consequence, the accuracy of the EGM2008 gravity field  
142 functionals varies over different parts of Earth. EGM2008 is most accurate (i.e., lowest  
143 commission errors) in regions with high-quality terrestrial gravity data sets (i.e., dense  
144 coverage, sufficient accuracy) available for its construction.

145 For practical applications, EGM2008-based functionals of the gravity field are obtained  
146 through harmonic synthesis of the model coefficients. Harmonic synthesis (e.g., Torge 2001,  
147 p. 271) can be accomplished, e.g., using the publicly available high-degree harmonic  
148 synthesis software `harmonic_synth` (Holmes and Pavlis 2008). The software is capable of  
149 computing a variety of EGM2008 gravity field functionals (e.g., geoid and quasigeoid  
150 heights, gravity anomalies and disturbances, vertical deflections), either in terms of scattered  
151 locations or points arranged as equidistant grids.


152 When using the `harmonic_synth` software, only the scattered point option allows for  
153 (individual) ellipsoidal heights of the topography, while grid computations are carried out at  
154 some given *constant* ellipsoidal height (e.g., surface of a reference ellipsoid). The EGM2008  
155 quasigeoid heights over Germany – computed with the `harmonic_synth` scattered point option  
156 at the ellipsoidal heights of the topography are shown in Fig. 1.



157

158 Fig. 1: EGM2008 quasigeoid heights over Germany (spectral degrees 2 to 2190, unit in  
 159 metres)

160



161

162 Fig. 2: EGM2008 commission errors over Germany (spectral degrees 2 to 2190, unit in  
 163 metres)

164 Users of EGM2008 also have the option of downloading pre-computed grids of EGM2008-  
 165 based functionals from the EGM2008 website ([http://earth-  
 166 info.nga.mil/GandG/wgs84/gravitymod/new\\_egm/TEST\\_RESULTS/results.html](http://earth-info.nga.mil/GandG/wgs84/gravitymod/new_egm/TEST_RESULTS/results.html) and  
 167 [http://earth-info.nga.mil/GandG/wgs84/  
 168 gravitymod/egm2008/index.html](http://earth-info.nga.mil/GandG/wgs84/gravitymod/egm2008/index.html)). These grids were computed at the surface of the reference ellipsoid using harmonic\_synth's grid mode. Hence,


169 they provide EGM2008 geoid and quasigeoid heights, gravity anomalies and vertical  
170 deflections at a constant ellipsoidal height (of 0 m) above the ellipsoid. The role of the  
171 ellipsoidal height used in the synthesis is further dealt with in Section 5.

172 Maps of EGM2008 commission errors were computed by the EGM2008 development team  
173 for quasi/geoid heights, gravity anomalies and vertical deflections using a dedicated error  
174 propagation technique described in Pavlis and Saleh (2004). For areas with rather scarce  
175 surface gravity coverage (for instance, parts of Africa, South America and Asia), commission  
176 errors for EGM2008 quasi/geoid undulations are estimated to be at the level of ~15cm with  
177 maximum uncertainties encountered in the mountainous parts of Asia and South America  
178 (around ~30-40 cm) and Antarctica (~100 cm). In contrast to this, the lowest commission  
179 errors are found over most parts of Europe, Oceania, North America and – because of the use  
180 of dense sets of altimetry-derived gravity – the oceans (see Pavlis et al. 2008). For those  
181 regions with high-quality surface gravity available, the EGM2008 quasi/geoid commission  
182 errors are mostly at the level of ~5 cm. A detailed map of the EGM2008 quasi/geoid  
183 commission errors over Germany is shown in Fig. 2, where the error estimates range from 3  
184 cm to 10 cm, with an average value of 5 cm. Section 5 will demonstrate that these ‘official’  
185 commission error estimates are rather pessimistic for Germany.

### 186 **3. Residual Terrain Modelling (RTM) approach**

187 The truncation of EGM2008 model coefficients at spherical harmonic degree 2190 produces  
188 an omission error (Torge 2001, p 273). In other words, the fine-structure of Earth’s gravity  
189 field at scales less than 5 arc minutes is not contained in the EGM2008-based gravity field  
190 functionals. As shown in Hirt (2010), residual terrain modelling (RTM) is one approach that  
191 is suited to compute and reduce this omission error. The basic idea of the RTM method  
192 (Forsberg 1984) is to construct residual elevations as the difference between a high-resolution  
193 elevation model of the topography and some long-wavelength ‘reference’ topography, which  
194 acts as a high-pass filter. The residual elevations are then used to compute RTM gravity field  
195 functionals, in order to reduce the omission error of the truncated EGM2008 model to some  
196 extent (Hirt 2010, Hirt et al. 2010a, 2010b). In the construction of EGM2008, a variant of the  
197 RTM technique was employed for the ‘prediction’ of band-limited gravity anomalies over  
198 areas with relatively poor gravity data coverage (Pavlis et al. 2007).


199



200

201 Fig. 3: RTM elevations (SRTM minus DTM2006.0) over Germany (unit in metres)

202



203

204 Fig. 4: RTM quasigeoid heights over Germany (unit in metres)

205 In this work, we use the 3 arc second SRTM (Shuttle Radar Topography Mission) data set  
206 (release 4.1 by Jarvis et al. 2008) as the high-resolution elevation model. The long-  
207 wavelength reference topography is provided by the spherical harmonic expansion of the  
208 DTM2006.0 data base (Pavlis et al. 2007, Saleh and Pavlis 2003), which is an associated  
209 EGM2008 product. Expanded to harmonic degree 2160, DTM2006.0 elevations ‘remove’ a  
210 large part of those gravity field signals from the SRTM topography, that are already implied  
211 by EGM2008 (Hirt 2010). The transformation of RTM elevations to RTM quasigeoid heights  
212 is accomplished using mass-density forward modelling (e.g., Torge 2001, p. 260, Forsberg  
213 1984, Nagy et al. 2000, Hirt et al. 2010a) with software based on the TC program (Forsberg  
214 1984) and a density assumption of constant mass-density of  $2670 \text{ kg/m}^3$ . The resulting RTM  
215 quasigeoid heights are the contribution of the RTM model topography to the EGM2008-  
216 omitted signals. Mainly because of short-scale (beyond EGM2008 resolution) mass-density  
217 anomalies in the real topography, the RTM approach only approximates the EGM2008 signal  
218 omission to some extent (Hirt 2010). To reduce the EGM2008 quasigeoid omission error,  
219 RTM quasigeoid heights are simply added to those from EGM2008 (Hirt et al. 2010b).

220 Fig. 3 shows the SRTM minus DTM2006.0 elevations over Germany and parts of the  
221 neighbouring countries. RTM quasigeoid heights were computed in terms of a high-  
222 resolution  $0.3' \times 0.3'$  grid (equivalent to a resolution of 550 m in latitude  $\times$  350 m in longitude)  
223 covering the whole of Germany (Fig. 4). Each RTM quasigeoid height originates from the  
224 evaluation of the SRTM-DTM2006.0 RTM data within 200 km radius around any  
225 computation point (extending the area shown in Fig. 3). Over the North Sea and Baltic Sea,  
226 DTM2006.0 and SRTM elevations were set to zero, so as to avoid artefacts coming from the  
227 bathymetry contained in DTM2006.0.

228 Over Germany, the RTM quasigeoid heights (Fig. 4) possess – on average – a signal strength  
229 of 1.3 cm (RMS, root mean square). In rugged terrain, such as the German Alps (South of  
230  $47.5^\circ$  latitude), the amplitudes of the RTM quasigeoid are larger with maximum values of  
231  $\sim 17$  cm, while the RTM approach fails to model the omission error of EGM2008 over level  
232 terrain (Figs. 3 and 4).

#### 233 **4. Comparison data sets**

234 As comparison data for an assessment of EGM2008 over Germany, this study utilizes  
235 quasigeoid heights (also denoted height anomalies) from the GCG05 quasigeoid model, from  
236 GPS/levelling and quasigeoid height differences from astronomical levelling. Because similar  
237 gravity data sets were likely used in the development of the EGM2008 and GCG05, these  
238 models are inevitably dependent to some extent. This is why GCG05 cannot be used for a  
239 truly independent assessment of EGM2008. Rather, the comparisons involving GCG05 are  
240 used to examine different EGM2008 evaluation variants, including RTM-based omission  
241 error corrections over a dense grid (Section 5). In contrast to GCG05, the GPS/levelling  
242 stations and astrogeodetic quasigeoid height differences are independent of EGM2008 and  
243 therefore a useful complement to the GCG05 comparisons. It should be noted that there exists  
244 a tight relation between the GPS/levelling set used here and GCG05 (described below).



#### 245 **4.1 The GCG05 quasigeoid model**

246 The GCG05 (German Combined Quasigeoid 2005) quasigeoid model is the official height  
247 reference surface of the AdV (Arbeitsgemeinschaft der Vermessungsverwaltungen der  
248 Länder) and can be used for the conversion between ellipsoidal and physical heights over  
249 Germany (BKG 2006, Liebsch et al. 2006). GCG05 provides 120,530 quasigeoid heights on a  
250 grid of 1.0'×1.5' (resolution of ~1.8 km in latitude × ~1.7 km in longitude). The accuracy of  
251 the GCG05 quasigeoid heights is specified to be 1-2 cm (BKG 2006). Locally, the accuracy  
252 of GCG05 quasigeoid height differences can be better than 1 cm (Hirt et al. 2007, Hirt et al.  
253 2008), see also Sect. 5.3.


254 GCG05 is a gravimetric quasigeoid model that originates from two independent RCR-  
255 computations performed at Leibniz Universität Hannover (Institut für Erdmessung) and  
256 Bundesamt für Kartographie und Geodäsie (BKG). According to Liebsch et al. (2006), the  
257 model is based on ~430,000 gravity anomalies, high-resolution elevation data and ~900  
258 GPS/levelling points. For the RCR-procedure, the EIGEN-CG01C global gravity field model  
259 (Reigber et al. 2006), expanded to degree 360, was used as reference. In addition to surface  
260 gravity data, this global model incorporates more than 2 years of CHAMP and 3.5 months of  
261 GRACE satellite gravity data (Reigber et al. 2006), conferring highly-accurate long- and  
262 medium-wavelength information to GCG05. The techniques used for the computation of the  
263 quasigeoid heights from the gravity anomalies are least-squares spectral combination  
264 (Leibniz Universität Hannover) and point mass adjustment (BKG). Both solutions were  
265 combined with GPS/levelling quasigeoid heights and arithmetically averaged to yield the  
266 GCG05 quasigeoid model (Schirmer et al. 2006, Liebsch et al. 2006).

#### 267 **4.2. Quasigeoid heights from GPS/levelling**

268 A set of GPS/levelling points (Ihde and Sacher 2002) was kindly made available by BKG.  
269 This data set provides quasigeoid heights as the differences between GPS-observed  
270 ellipsoidal heights and spirit-levelled normal heights at 675 locations scattered over  
271 Germany. The GPS/levelling quasigeoid heights are independent of EGM2008 and can be  
272 assumed to be accurate to a few cm. This set was also used by Gruber (2009) for an  
273 evaluation of EGM2008 (without RTM augmentation).

274 The GCG05 model and the quasigeoid heights at the 675 GPS/levelling points are tightly  
275 related, but not identical, as explained next. The 675 GPS/levelling points (Ihde and Sacher  
276 2002) form a subset of the ~900 GPS/levelling points (Liebsch et al. 2006), but were not  
277 directly used in the construction of the GCG05 model. Prior to the construction of GCG05,  
278 the ellipsoidal heights of the 675 GPS/levelling points were adapted to ETRS89 (European  
279 Terrestrial Reference System 89), as realised by the SAPOS (Satellitenpositionierungsdienst  
280 der deutschen Landesvermessungen) reference station network. The adaption of ellipsoidal  
281 heights was done in most different ways by the state survey agencies of Germany (Liebsch et  
282 al 2006, p. 135). Hence, the quasigeoid heights are different in both GPS/levelling sets (see  
283 also Liebsch et al. 2006 p. 136). As an immediate consequence, there exist small differences  
284 between the GCG05 quasigeoid heights and those of the 675 GPS/levelling points (Ihde and

285 Sacher 2002). Fig. 5 shows that these differences are at the level of a few cm  
286 (min/max/mean/rms: -5.3/9.1/0.5/1.5 cm), see also Liebsch et al. (2006), p. 136.




287

288 Fig. 5: Differences between quasigeoid heights of the 675 GPS/levelling points and GCG05  
289 (unit in metres)

### 290 4.3. Astrogeodetic quasigeoid differences

291 Finally, this study uses two local profiles of highly-accurate quasigeoid height differences  
292 that were computed from astrogeodetic vertical deflections (Fig. 6). The vertical deflections  
293 were observed using the Hannover digital zenith camera (Hirt et al. 2010c) at densely-spaced  
294 stations. The first profile of 114 observed astrogeodetic stations over a distance of 63 km  
295 length (Fig. 6A) crosses the Harz Mountains in Northern Germany (Hirt et al. 2008). The  
296 second profile (Fig. 6B) is located in the Isar Valley, Bavaria, has a length of 23 km and  
297 consists of 103 observations (Hirt et al. 2007, Hirt and Flury 2008). In both test areas, the  
298 astrogeodetic vertical deflections were interpolated utilizing high-resolution elevation data  
299 and transformed to quasigeoid height differences by means of Helmert's path integral (see  
300 Hirt and Flury 2008).

301 The accuracy of the astrogeodetic quasigeoid height differences was estimated to be 1-2 mm  
302 over the length of both profiles (Hirt et al. 2008, Hirt and Flury 2008). This makes both data  
303 sets well-suited for the local validation of EGM2008. It should be noted that both profiles  
304 were connected with additional vertical deflection observations to form a ~600 km North-  
305 South profile (Voigt et al. 2008, 2009). This data set was used for regional comparisons with  
306 EGM2008 (Berichte 2010, and Ihde et al. 2010), however without the omission error  
307 modelling as is done here.



308

309 Fig. 6: Location of the astrogeodetic quasigeoid profiles. A: Harz Mountains profile, B: Isar  
 310 Valley profile. The background topography are SRTM heights in metres.

## 311 5. Comparisons

### 312 5.1 EGM2008 vs. GCG05

313 The zero-tide<sup>1</sup> version of EGM2008 was evaluated with the scattered point option of the  
 314 harmonic\_synth software (Holmes and Pavlis 2008) over the spherical harmonic band from  
 315 degree 2 to 2190 at the geodetic coordinates latitude and longitude of the 120,530 GCG05  
 316 grid points. As a first processing variant, a constant ellipsoidal height of 0 m (i.e., surface of  
 317 the reference ellipsoid) was used. This replicates the case of using pre-calculated grids from  
 318 the EGM2008 website. As a second processing variant, ellipsoidal heights of the topography  
 319 were ‘constructed’ as the sum of SRTM elevations (in approximation, these are heights above  
 320 mean sea level) and GCG05 quasigeoid heights and subsequently used in the synthesis  
 321 procedure (cf. Claessens et al. 2009).


322 RTM quasigeoid heights were obtained for at the GCG05 grid points through interpolation of  
 323 the 0.3'×0.3' RTM quasigeoid grid (Fig. 3). The two synthesis variants and the optional  
 324 consideration of RTM effects allow four different comparisons between GCG05 and  
 325 EGM2008 (Fig. 7). The descriptive statistics of the differences are reported in Tab. 1. In each  
 326 of the comparisons, the mean value of the differences was subtracted (known as 1-parameter  
 327 or bias-fit) to eliminate the impact of different vertical datums (zero levels) and very long  
 328 wavelength errors of the data sets (cf. Featherstone 2001 and Ihde et al. 2010).

---

<sup>1</sup> Zero-tide means that the lunisolar permanent deformation of the Earth is included while the attraction effect is eliminated (Torge 2001, p. 77). The use of the zero-tide system follows a recommendation of the International Association of Geodesy (IAG) and is the preferred tide-system in practical quasi/geoid computations (e.g., Denker et al. 2009, Featherstone et al. 2010). Unfortunately, GCG05 cannot be considered a pure zero-tide model (Liebsch 2011, pers. comm.) which may cause small discrepancies in the comparisons. A detail discussion and analysis of the tide systems of the GCG05 input data sets is beyond the scope of the present study.

329 The comparison between GCG05 and EGM2008 evaluated at the ellipsoidal height = 0 m  
 330 (Fig. 7A) shows RMS errors of 3.3 cm with maximum discrepancies of ~25 cm occurring in  
 331 the German Alps, South of ~48°N. Evaluation of EGM2008 quasigeoid height at the  
 332 ellipsoidal height of the topography (Fig. 7B) improves the agreement with GCG05 in the  
 333 elevated or mountainous parts of Germany by ~5-10 cm. This is seen for the Harz Mountains  
 334 (51.7°N, 10.5°E), the Black Forest (47.8°N, 8°E), and over wide areas of Bavaria. The largest  
 335 improvement of up to ~20 cm is found over the German Alps.

336



337

338 Fig. 7: Differences between the German Quasigeoid model GCG05 and variants of  
 339 EGM2008. A: GCG05–EGM2008 (evaluated on the ellipsoid,  $h = 0$ ), B: GCG05–EGM2008

340 (evaluated at the ellipsoidal height of the topography), C: GCG05–[EGM2008 (evaluated on  
 341 the ellipsoid,  $h = 0$ ) + RTM], D: GCG05–[EGM2008 (evaluated at the ellipsoidal height of  
 342 the topography) + RTM]. Units in metres.

343

344 Tab. 1: Descriptive statistics of the quasigeoid differences between GCG05 and EGM2008  
 345 variants (bias-fit, 120,530 points)

Comparison	Min [cm]	Max [cm]	RMS [cm]
GCG05 – (EGM2008, $h=0$ )	-24.7	24.1	3.3
GCG05 – (EGM2008, $h=topo$ )	-19.1	23.5	3.2
GCG05 – [(EGM2008, $h=0$ ) + RTM ]	-22.1	11.0	3.1
GCG05 – [(EGM2008, $h= topo$ ) + RTM]	-17.4	10.3	3.0

346

347 Additional consideration of RTM quasigeoid heights in the comparisons (Figs. 7C and 7D)  
 348 reduces most of the short-wavelength (scales of  $\sim 10$  km and below) error patterns seen  
 349 previously. The most striking example are for the German Alps (see also Hirt et al. 2010a),  
 350 but also for many other regions of Germany, except for the parts of Northern Germany with  
 351 lower relief. The best model fit is observed for EGM2008 evaluated at the topography with  
 352 the support of RTM quasigeoid heights beyond EGM2008’s resolution (Fig. 7D). For this  
 353 variant, the maximum differences are significantly reduced with respect to EGM2008-only  
 354 (cf. Tab. 1), while the RMS differences only slightly improve to 3 cm. This behaviour is  
 355 attributed to the medium-wavelength difference patterns (scales of  $\sim 100$ -200 km and larger),  
 356 that are present in each of the four comparisons (Fig. 7) and are discussed in Section 5.2.

357 It should be noted that small-amplitude high-frequency difference patterns remain even in  
 358 Fig. 7D where RTM quasigeoid heights were used to augment the EGM2008 resolution.  
 359 These high-frequency effects which occur with wavelengths of  $\sim 20$  km are further analysed  
 360 in Section 5.3.

361 Given that the 3 cm RMS value reflects the commission errors of GCG05, EGM2008 and  
 362 RTM quasigeoid heights, a good quality of the three data sets is indicated. However, GCG05  
 363 quasigeoid does not allow a truly independent validation of EGM2008 (see Sect. 4).  
 364 Nonetheless, the comparisons provide a good feedback on the different gravimetric modelling  
 365 strategies employed (harmonic analysis in case of EGM2008, and spectral combination/point  
 366 mass adjustment for GCG05).


## 367 5.2 EGM2008 vs. GPS/levelling

368 For the comparisons with the GPS/levelling quasigeoid heights, EGM2008 and RTM  
 369 quasigeoid heights were evaluated in the fashion described above. However, due to the  
 370 precise ellipsoidal heights provided by the GPS component, a ‘construction’ of ellipsoidal  
 371 heights was not necessary for the synthesis task. The descriptive statistics of the four different  
 372 comparison variants among EGM2008 and the GPS/levelling set is given in Tab. 2. The

373 difference patterns between GPS/levelling quasigeoid heights and the two selected variants  
 374 ‘EGM2008 (evaluated at height = 0 m)’ and ‘EGM2008 (evaluated at the topography) +  
 375 RTM augmentation’ are shown in Fig. 8. The difference patterns and behaviour of the  
 376 statistics are fairly comparable to the GCG05 comparisons before (compare Tab. 1 and 2),  
 377 which was expected due to the tight relation between the GPS/levelling set and the GCG05  
 378 model. Again, inclusion of RTM quasigeoid heights leads to a significant reduction in the  
 379 extreme discrepancies, while there is only a small improvement in the RMS, from 3.6 cm to  
 380 3.3 cm.

381 Using the same GPS/levelling data set, Gruber (2010) found a similar RMS value (3.8 cm)  
 382 for the EGM2008-only comparisons. For the GCG05 GPS/levelling set of ~900 points, Ihde  
 383 et al. (2010) published a RMS value of 3.0 cm, reflecting the better quality of their newer  
 384 GPS data.

385



386

387 Fig. 8: Differences between GPS/levelling quasigeoid heights and variants of EGM2008. A:  
 388 GPS/levelling – EGM2008 (evaluated on the ellipsoid), B: GPS/levelling – [EGM2008  
 389 (evaluated at the ellipsoidal height of the topography) + RTM]. Units in metres.

390 Tab. 2: Descriptive statistics of the quasigeoid differences between GPS/levelling and  
 391 EGM2008 variants (bias-fit, 675 points)

Comparison	Min [cm]	Max [cm]	RMS [cm]
GPS/lev – (EGM2008, h=0)	-9.6	21.8	3.6
GPS/lev – (EGM2008, h= topo)	-10.5	18.0	3.6
GPS/lev – [(EGM2008, h=0) + RTM]	-8.8	13.3	3.5

GPS/lev – [(EGM2008, h= topo) + RTM]	-9.7	10.0	3.3
--------------------------------------	------	------	-----

392

393 The comparison between EGM2008 and the truly-independent GPS/levelling heights exhibits  
 394 medium-wavelength error patterns with coarsely 5 cm amplitude (e.g., yellow areas over  
 395 Bavaria and Thuringia) which were similarly seen before in the GCG05 comparisons. Some  
 396 correlations can be observed between the commission error map (Fig. 2) and the difference  
 397 patterns in Fig. 7D. This, together with the ‘official’ accuracy estimates of the data sets in  
 398 mind (Sect. 3 and 4), would suggest that these patterns reflect EGM2008 commission errors  
 399 rather than those of the comparison data. However, there exist at least two further sources of  
 400 error which may explain parts of the medium-wavelength error patterns.


401 First, inhomogenities in the GPS/levelling data, particularly in the ellipsoidal GPS heights,  
 402 might be responsible, occurring as ‘step-like effect’ at some of the state boundaries, e.g.  
 403 between Bavaria and Thuringia (Voigt et al. 2009; Feldmann-Westendorff 2010, pers.  
 404 comm.). This effect might have propagated into the GPS/levelling set and GCG05 model,  
 405 and, in turn, into the differences seen in Fig. 7 and 8. Within the framework of the current  
 406 renewal of the German first-order levelling network (Deutsches Haupthöhennetz DHHN, e.g.,  
 407 Jahn 2010, Feldmann-Westendorff 2009, Feldmann-Westendorff and Jahn 2006), 250 high-  
 408 quality GNSS/levelling stations (and 100 absolute gravity stations) will become available in  
 409 the near future. The quasigeoid heights of the GNSS/levelling stations are based on state-of-  
 410 the-art GNSS measurements and a re-observation of the first-order levelling lines. Owing to  
 411 the expected level of accuracy (1 cm and better), this data set is likely to be suited for a future  
 412 investigation of EGM2008 (and GCG05) commission errors in particular, and, of course, for  
 413 evaluating future geopotential models in general.

414 Second, also the EIGEN-CG01C geopotential model used in the GCG05 construction might  
 415 be a possible explanation for parts of the medium-wavelength differences in Fig. 7 (see also  
 416 Ihde et al. 2010). However, because the difference patterns between EGM2008 and the  
 417 GPS/levelling data (Fig. 8) are independent of the EIGEN-model, it can be concluded that  
 418 EIGEN-CG01C is not the main contributor to the discrepancies. High-accuracy geopotential  
 419 models, which are currently constructed based on the GOCE satellite gravity field mission  
 420 (e.g., Rummel et al. 2009) can be expected to allow clarification of the medium-wavelength  
 421 difference patterns. This is because of the expected geoid cm-accuracy over scales of ~100  
 422 km (e.g., Rummel 2005) and, importantly, because of the fact that the GOCE observations  
 423 are independent of all data sets involved here (specifically EGM2008, GCG05 and the  
 424 underlying geopotential models).

### 425 **5.3 EGM2008 vs. astronomical levelling**

426 Finally, the residuals between the highly-accurate astrogeodetic quasigeoid height differences  
 427 and the four EGM2008 variants are shown in Fig. 9A for the Harz Mountains profile (Fig.  
 428 6A) and Fig. 9B for the Isar Valley profile (Fig. 6B). The descriptive statistics of the  
 429 comparisons are given in Tab. 3 and 4. For the sake of completeness, the residuals with  
 430 respect to GCG05 (cf. Hirt et al. 2007, 2008) are also displayed in Fig. 9, showing the very

431 good agreement (better than cm-level) between the independent astrogeodetic and  
 432 gravimetric quasigeoid solutions at local scales.



433  
 434 Fig. 9: Differences between the astrogeodetic quasigeoid heights and the EGM2008 variants  
 435 (red, blue lines) and differences between the astrogeodetic quasigeoid heights and GCG05  
 436 (black lines). A: Harz Mountains, B: Isar Valley. The quasigeoid heights of the first profile  
 437 points were set to zero for all data sets.

438 Tab. 3: Harz Mountains: Descriptive statistics of the quasigeoid differences between the  
 439 astrogeodetic solution and EGM2008 variants (first station set to zero for any data set, 114  
 440 points)

Comparison	Min [cm]	Max [cm]	Mean [cm]	RMS [cm]
Astro – (EGM2008, h=0)	-9.0	0.3	-4.2	5.1
Astro – (EGM2008, h= topo)	-1.7	1.8	-0.0	1.1
Astro – [(EGM2008, h=0) + RTM]	-8.7	0.0	-3.7	4.8
Astro – [(EGM2008, h= topo) + RTM]	-1.2	2.4	0.5	1.0
Astro – GCG05	-0.6	0.7	0.1	0.3

441  
 442 Tab. 4: Isar Valley: Descriptive statistics of the quasigeoid differences between the  
 443 astrogeodetic solution and EGM2008 variants (first station set to zero for any data set, 103  
 444 points)

Comparison	Min [cm]	Max [cm]	Mean [cm]	RMS [cm]
Astro – (EGM2008, h=0)	-11.5	0.0	-5.5	6.1
Astro – (EGM2008, h= topo)	-8.6	0.0	-4.4	4.8
Astro – [(EGM2008, h=0) + RTM]	-6.1	0.2	-2.5	3.0
Astro – [(EGM2008, h= topo) + RTM]	-3.2	0.0	-1.4	1.6
Astro – GCG05	-1.5	1.3	-0.1	0.8



446 Fig. 9 demonstrates the effect of not evaluating EGM2008 at the topography (solid versus  
447 dotted lines). For both profiles, EGM2008, evaluated at the ellipsoidal height of the  
448 topography and augmented by RTM, produces the lowest RMS discrepancies of 1.0 cm  
449 (Harz) and 1.6 cm (Isar Valley). These comparisons show that EGM2008 – over well-  
450 surveyed areas and with augmentation of RTM data in mountainous terrain – is capable of  
451 delivering differences of quasigeoid heights near or at the cm-level. For other parts of  
452 Germany, this finding is corroborated by the structure of the difference patterns of the  
453 EGM2008/RTM comparisons with GCG05 and GPS/levelling (Figs. 7D and 8B). Over many  
454 regions, for instance large parts of Bavaria, the differences patterns are fairly constant, so  
455 would cancel out to some extent when quasigeoid height *differences* are computed from  
456 EGM2008. This demonstrates EGM2008 can be a source of quasigeoid height differences  
457 near the cm-level at local scales, over distances of few tens of km. However, GCG05 is an  
458 even more accurate source for quasigeoid height differences over Germany (Fig. 9).

459 For the comparisons involving EGM2008, oscillating differences with roughly ~20 km  
460 wavelength (i.e., the resolution of EGM2008) and amplitudes of ~2 cm are visible in Fig. 9.  
461 Similar high-frequency difference patterns occurred previously in the comparisons between  
462 GCG05 and EGM2008/RTM (Fig. 7D). This, together with the sub-cm agreement between  
463 GCG05 and the astrogeodetic solutions (Fig. 9) provides some evidence that the small high-  
464 frequency error patterns, as visible over parts of Germany (Fig. 7D) does not originate from  
465 GCG05, but from EGM2008 or from the RTM omission error corrections.

## 466 **6. Conclusions**

467 The present study evaluated the EGM2008 global geopotential model over Germany as an  
468 example region where dense gravity data sets were used for the model's development. For the  
469 EGM2008 evaluation, quasigeoid heights or quasigeoid height differences sourced from three  
470 different terrestrial data sets were used. To improve upon the short-wavelength signals,  
471 EGM2008 was augmented by quasigeoid heights from residual terrain model data. In  
472 elevated or mountainous terrain, this is efficient to reduce the omission error of the  
473 EGM2008 quasigeoid heights. The discrepancies with respect to GCG05 and GPS/levelling  
474 quasigeoid heights are at the level of 3 cm. Locally, say over distances of a few tens of km,  
475 EGM2008 (augmented by RTM) may deliver quasigeoid height differences near or at the cm-  
476 level, as was indicated by the comparisons with astrogeodetic data.

477 The comparisons involving the astrogeodetic data provide evidence that EGM2008, though  
478 being a good model over Germany, does not yet reach the quality of the GCG05 national  
479 quasigeoid model for quasigeoid height differences. The comparisons between EGM2008  
480 and the three quasigeoid data sets show that the official EGM2008 commission error  
481 estimates (~5 cm for Germany) are too pessimistic. The medium-wavelength error patterns,  
482 which became visible in the comparisons between EGM2008 and GCG05 and EGM2008 and  
483 GPS/levelling could not be unambiguously attributed to one (or more) of the models used in  
484 this study. However, new data sets (quasigeoid heights from the DHHN renewal and from the  
485 GOCE mission) are expected to yield further insight into the discrepancies between  
486 EGM2008, GCG05 and GPS/levelling.

487 The different evaluation variants of EGM2008 used in this study have underlined the  
488 importance of using ellipsoidal heights of the topography in the synthesis of gravity field  
489 functionals. Evaluation at the ellipsoidal surface (ellipsoidal height = 0 m) may contaminate  
490 the computed quasigeoid heights by ~5-20 cm in elevated and mountainous areas of  
491 Germany. If EGM2008 is used for the prediction of quasigeoid heights or other functionals  
492 (e.g., gravity, vertical deflections) at the Earth's surface, some care should be exercised with  
493 pre-computed grids of EGM2008 functionals (and the use of the harmonic\_synth software in  
494 grid mode), unless the influence of the topography is corrected otherwise.

495 As a general conclusion, the results of this study show the high quality of EGM2008 over  
496 densely surveyed regions and confirms the advancements made in global gravity field  
497 modelling, as demonstrated by development of EGM2008. While a similarly good quality is  
498 expected or indicated for other well-surveyed regions (see Newton's Bulletin 2009), it  
499 should be noted that EGM2008 commission errors may be significantly higher in areas of  
500 poor gravity data coverage (cf. Pavlis et al. 2008).

501 **Acknowledgements** The author acknowledges funding through Australian Research Council  
502 grant DP0663020. Thanks go to the EGM2008 Development Team for making their model  
503 freely available. The GCG05 model and GPS/levelling points were provided by BKG (Uwe  
504 Schirmer). This support is kindly acknowledged. Valuable comments on the manuscript by  
505 Kevin Fleming and by two reviewers, particularly reviewer #2, are appreciated.

## 506 **References**

- 507  
508 Andersen O.B., Knudsen P. and Berry P.A.M. (2010): The DNSC08GRA global marine gravity field from  
509 double retracked satellite altimetry, *J. Geod.*, 84(3), pp 191-199, DOI: 10.1007/s00190-009-0355-9.
- 510 Berichte (2010): Berichte 2009 aus der Fachrichtung Geodäsie und Geoinformatik, Februar 2010, Folge 60.  
511 Leibniz Universität Hannover.
- 512 BKG (2006): Quasigeoid der Bundesrepublik Deutschland. Die Höhenreferenzfläche der Arbeitsgemeinschaft  
513 der Vermessungsverwaltungen der Länder GCG05 (German Combined Quasigeoid 2005). Geodätische  
514 Basisdaten Bundesrepublik Deutschland. URL: [www.geodatenzentrum.de/docpdf/quasigeoid.pdf](http://www.geodatenzentrum.de/docpdf/quasigeoid.pdf).
- 515 Claessens S.J., Hirt C., Amos M.J., Featherstone W.E. and Kirby J.F. (2011): The NZGEOID09 New Zealand  
516 quasigeoid model, *Survey Review*, 43(319), pp 2-15.
- 517 Claessens S.J., Featherstone W.E., Anjasmara I.M., Filmer M.S. (2009): Is Australian data really validating  
518 EGM2008 or is EGM2008 just in/validating Australian data, In: *Newton's Bulletin* (2009), pp 207-251.
- 519 Denker H., Barriot J.-P., Barzaghi R., Fairhead, D., Forsberg R., Ihde J., Kenyars A., Marti U., Sarrailh M., and  
520 Tziavos I.N. (2009): The development of the European gravimetric geoid model EGG07. in: Sideris M.G  
521 (ed) *Observing Our Changing Earth*, Springer, Berlin Heidelberg New York, pp 177-186.
- 522 EGM Development Team (2008): Earth Gravitational Model 2008 – readme files. URL: [http://earth-](http://earth-info.nga.mil/GandG/wgs84/gravitymod/egm2008/)  
523 [info.nga.mil/GandG/wgs84/gravitymod/egm2008/](http://earth-info.nga.mil/GandG/wgs84/gravitymod/egm2008/).
- 524 Featherstone W.E. (2001): Absolute and relative testing of gravimetric geoid models using Global Positioning  
525 System and orthometric height data, *Computers and Geosciences*, 27(7), pp 807-814, DOI:  
526 10.1016/S0098-3004(00)00169-2.
- 527 Featherstone W.E., Kirby J.F., Hirt C., Filmer M.S., Claessens S.J., Brown N.J., Hu G. and Johnston G.M. (in  
528 press): The AUSGeoid09 model of the Australian Height Datum, *Journal of Geodesy*, accepted (Oct  
529 2010), DOI: 10.1007/s00190-010-0422-2.
- 530 Feldmann-Westendorff U. and Jahn C.-H. (2006): GNSS-Höhenbestimmung in einem einheitlichen  
531 Raumbezug. 66. DVW-Seminar "GPS und GALILEO", Deutscher Verein für Vermessungswesen e. V. -  
532 Gesellschaft für Geodäsie, Geoinformation und Landmanagement, 21.-22.02.2006, Darmstadt,  
533 Schriftenreihe 49/2006, pp 147-171.

- 534 Feldmann-Westendorff U. (2009): Von der See bis zu den Alpen: Die GNSS-Kampagne 2008 im DHHN 2006-  
535 2011. DVW Seminar "GNSS 2009: Systeme, Dienste und Anwendungen, 18/19.03.2009, TU Dresden.  
536 DVW Schriftenreihe Nr 57, pp. 95-112.
- 537 Forsberg R. (1984): A study of terrain reductions, density anomalies and geophysical inversion methods in  
538 gravity field modelling. Report 355, Department of Geodetic Science and Surveying, Ohio State  
539 University, Columbus.
- 540 Förste C., Stubenvoll R., König R., Raimondo J.-C., Flechtner F., Barthelmes F., Kusche J., Dahle C.,  
541 Neumayer H., Biancale R., Lemoine J.-M. and Bruinsma S. (2009): Evaluation of EGM2008 by  
542 comparison with other recent global gravity field models. In: Newton's Bulletin (2009), pp 26-37.
- 543 Gruber, T. (2009): Evaluation of the EGM2008 Gravity Field by Means of GPS Levelling and Sea Surface  
544 Topography Solutions, In: Newton's Bulletin (2009), pp 3-17.
- 545 Hirt C., Denker H., Flury J., Lindau A. and Seeber G. (2007): Astrogeodetic Validation of Gravimetric  
546 Quasigeoid Models in the German Alps - First Results, Proceed. of the 1st International Symposium of  
547 the International Gravity Field Service (IGFS), Istanbul, Turkey, Harita Dergisi, Special Issue 18, pp 84-  
548 89.
- 549 Hirt C., Feldmann-Westendorff U., Denker H., Flury J., Jahn C.-H., Lindau A., Seeber G. and Voigt C. (2008):  
550 Hochpräzise Bestimmung eines astrogeodätischen Quasigeoidprofils im Harz für die Validierung des  
551 Quasigeoidmodells GCG05, Zeits. f. Verm u. Geoinf. (zfv) 133, pp 108-119.
- 552 Hirt C. and Flury J. (2008): Astronomical-topographic levelling using high-precision astrogeodetic vertical  
553 deflections and digital terrain model data. Journal of Geodesy 82(4-5), pp 231-248,  
554 DOI:10.1007/s00190-007-0173.
- 555 Hirt C. (2010): Prediction of vertical deflections from high-degree spherical harmonic synthesis and residual  
556 terrain model data. Journal of Geodesy 84(3), pp 179-190, DOI 10.1007/s00190-009-0354-x.
- 557 Hirt C., Marti U., Bürki B. and Featherstone W.E. (2010a): Assessment of EGM2008 in Europe using accurate  
558 astrogeodetic vertical deflections and omission error estimates from SRTM/DTM2006.0 residual terrain  
559 model data. Journal of Geophysical Research – Solid Earth, 115(B10): B10404, DOI:  
560 10.1029/2009JB007057.
- 561 Hirt C., Featherstone W.E and Marti U. (2010b): Combining EGM2008 and SRTM/DTM2006.0 residual terrain  
562 model data to improve quasigeoid computations in mountainous areas devoid of gravity data, Journal of  
563 Geodesy 84(9), pp 557-567, DOI 10.1007/s00190-010-0395-1.
- 564 Hirt C., Bürki B., Somieski A. and Seeber G. (2010c): Modern determination of vertical deflections using digital  
565 zenith cameras. J. Surv. Eng, Issue February 2010, pp 1-12, DOI: 10.1061/\_ASCE\_SU.1943-  
566 5428.0000009.
- 567 Holmes, S.A. and Pavlis N.K. (2007): Some aspects of harmonic analysis of data gridded on the ellipsoid.  
568 Proceed. of the 1st International Symposium of the International Gravity Field Service (IGFS), Istanbul,  
569 Turkey, Harita Dergisi, Special Issue 18, pp 151-156.
- 570 Holmes, S.A. and Pavlis N.K. (2008): Spherical harmonic synthesis software harmonic\_synth. Available online  
571 at: [http://earth-info.nga.mil/GandG/wgs84/gravitymod/new\\_egm/new\\_egm.html](http://earth-info.nga.mil/GandG/wgs84/gravitymod/new_egm/new_egm.html).
- 572 Ihde J. and Sacher M. (2002): EUREF Publikation 11/I, Mitteilungen des Bundesamtes für Kartographie und  
573 Geodäsie, Band 25, Frankfurt/Main, ISBN 3-89888-869-X.
- 574 Ihde, J., Wilmes H., Müller J., Denker H., Voigt C., and Hosse M. (2010): Validation of Satellite Gravity Field  
575 Models by Regional Terrestrial Data sets. In: System Earth via Geodetic-Geophysical Space Techniques  
576 Advanced Technologies in Earth Sciences, Part 3, pp 277-296.
- 577 Jarvis A., Reuter H.I., Nelson A. and Guevara E. (2008): Hole-filled SRTM for the globe Version 4, Available  
578 from the CGIAR-SXI SRTM 90m database: <http://srtm.csi.cgiar.org>.
- 579 Jahn C.-H. (2010): Das DHHN-Projekt: Ein Weg zum integrierten Raumbezug. Presented at Intergeo 7. Oct.  
580 2010, Cologne.
- 581 Jekeli C. (1988): The exact transformation between ellipsoidal and spherical expansions. Manuscripta Geodetica  
582 13(2), pp 106-133.
- 583 Jekeli C., Yanh H.J. and Kwon J.H. (2009): Evaluation of EGM08 – globally and locally in South Korea. In:  
584 Newton's Bulletin (2009), pp 39-49.

585 Kenyon S., Factor J., Pavlis N. and Holmes S. (2007): Towards the next Earth Gravitational Model. Paper  
586 presented at the Society of Exploration Geophysicists 77<sup>th</sup> Annual Meeting 23.-28 Sept. 2007, San  
587 Antonio, Texas.

588 Liebsch G., Schirmer U., Ihde J., Denker H., and Müller J. (2006): Quasigeoidbestimmung für Deutschland.  
589 Deutscher Verein für Vermessungswesen - Schriftenreihe, 49, pp 127-146.

590 Mayer-Gürr T. (2007): ITG-Grace03s: The latest GRACE gravity field solution computed in Bonn, Joint Int.  
591 GSTM and SPP Symposium, 15-17 Oct. 2007, Potsdam, Germany, [http://www.geod.uni-bonn.de/itg-](http://www.geod.uni-bonn.de/itg-grace03.html)  
592 [grace03.html](http://www.geod.uni-bonn.de/itg-grace03.html)

593 Nagy D., Papp G., and Benedek J. (2000): The Gravitational Potential and its Derivatives for the Prism J. Geod.,  
594 74(7-8), pp 552-560, DOI: 10.1007/s001900000116. Erratum in J. Geod. 76(8), pp 475-475, DOI:  
595 10.1007/s00190-002-0264-7.

596 Newton's Bulletin (2009): Newton's Bulletin Issue n° 4, April 2009 ISSN 1810-8555, Publication of the  
597 International Association of Geodesy and International Gravity Field Service.

598 Pavlis N.K. and Saleh J. (2004): Error Propagation with Geographic Specificity for Very High Degree  
599 Geopotential Models, Proceed. GGSM 2004 IAG International Symposium Porto, Portugal (ed. C. Jekeli  
600 et al.), Springer, Heidelberg, pp 149-154.

601 Pavlis N.K., Holmes S.A., Kenyon S.C., Schmidt D. and Trimmer R. (2004): A Preliminary Gravitational  
602 Model to Degree 2160. Proceed. GGSM 2004 IAG International Symposium Porto, Portugal (ed. C.  
603 Jekeli et al.), Springer, Heidelberg, pp 18-23.

604 Pavlis N.K., Factor J.K. and Holmes S.A. (2007): Terrain-related gravimetric quantities computed for the next  
605 EGM, Proceed. of the 1st International Symposium of the International Gravity Field Service (IGFS),  
606 Istanbul, Turkey, Harita Dergisi, Special Issue 18, pp 318-323.

607 Pavlis N.K., Holmes S.A., Kenyon S.C. and Factor J.K. (2008): An Earth Gravitational Model to Degree 2160:  
608 EGM2008, Presented at the 2008 General Assembly of the European Geoscience Union, Vienna, Austria,  
609 April 13-18, 2008.

610 Reigber C., Schwintzer P., Stubenvoll R., Schmidt R., Flechtner F., Meyer U., König R. Neumayer H., Förste  
611 C., Barthelmes F., Zhu S.Y., Balmino G., Biancale R., Lemoine J.-M., Meixner H. and Raimondo J.C.  
612 (2006): A High Resolution Global Gravity Field Model Combining CHAMP and GRACE Satellite  
613 Mission and Surface Data: EIGEN-CG01C. Scientific Technical Report STR06/07,  
614 GeoForschungsZentrum Potsdam.

615 Roman D.R., Wang Y.M. and Saleh J. (2010): Geodesy, Geoids, and Vertical Datums: A Perspective from the  
616 U.S. National Geodetic Survey. Paper presented at FIG Congress 2010, Technical Session 1C, 11-16.  
617 April 2010, Sydney, Australia.

618 Rummel R. (2005): Geoid and gravity in earth sciences - an overview. Workshop on Enabling Observation  
619 Techniques for Future Solid Earth Missions, Jan 30-Feb 01, 2003, Int Space Sci Inst Bern Switzerland,  
620 Earth Moon and Planets 94(1-2), pp 3-11.

621 Rummel R., Gruber T., Flury J. and Schlicht A. (2009): ESA's Gravity Field and Steady-State Ocean Circulation  
622 Explorer GOCE. Zeits. f. Verm u. Geoinf. (zfv) 3/2009, pp 125-130.

623 Saleh, J. and Pavlis N.K. (2003): The development and evaluation of the global digital terrain model DTM2002,  
624 Proceed. of the 3rd Meeting of the International Gravity and Geoid Commission of the International  
625 Association of Geodesy, Thessaloniki (ed. I. Tziavos), Editions Ziti, pp 207-212.

626 Schirmer U., Denker H., Ihde J., Liebsch G. and Müller J. (2006): A new combined height reference surface for  
627 Germany (GCG05). Presented at EUREF Symposium, Riga, Latvia 14 -17 Juni 2006.

628 Torge W. (2001): Geodesy. 3rd Edition, de Gruyter, Berlin, New York.

629 Voigt C., Denker H. and Hirt C. (2008): A GOCE regional validation experiment with vertical deflections in  
630 Germany. Presented at IAG Symposium on Gravity, Geoid and Earth Observation 23.-27. June 2008,  
631 Crete, Greece.

632 Voigt C., Denker H. and Hirt C. (2009): Regional astrogeodetic validation of GPS/levelling data and quasigeoid  
633 models. in: Sideris M.G (ed) Observing Our Changing Earth, Springer, Berlin Heidelberg New York, pp  
634 413-420.

635